THE EFFECT OF P-NITROCHLOROBENZENE ON HOMEOSTASIS QUANTITATIVE PARAMETERS OF KARST CAVE CLAYS AND ECUADOR SOILS MICROBIAL COMMUNITIES

In this paper it was given the effect of p-nitrochlorobenzene (NCB) on the homeostasis quantitative parameters of cave clays microbial communities from Western Ukraine and Abkhazia (Mushkarova Yama, Kuybushevskaya) and soils of Ecuador tropical ecosystems. For these microbial communities were determined maximum permissible concentrations and types of responses on xenobiotic. Microbial communities of Mushkarova Yama cave clays and rainforest soils of Ecuador were characterized by the first type of response. Microbial communities of Kuybushevskaya clays and mountain jungles of Ecuador were characterized by the second type of response. Maximum permissible concentration of NCB for Mushkarova Yama was 200 mg/l, for the other studied microbial communities – 300 mg/l. It was shown, that microbial communities were not only highly resistant to NCB but also interacted with it by destroying this xenobiotic and decreasing its concentration in 4 times.

Key words: p-nitrochlorobenzene (NCB), karst caves, Ecuador tropical ecosystems, microbial communities.

One of the main sources of pollution is waste waters from organic synthesis enterprises. There is special danger to the biosphere from xenobiotics included both nitro, chlorine and aromatic ring. P-nitrochlorobenzene (NCB) is one of such compounds. Its toxicity is determined by the aromatic ring and attached chlorine and nitro-group in the para-position. Bactericidal concentration for most soil and aquatic chemoorganotrophic microorganisms is 10 mg/l [1]. However, in several papers [2,3,4] has been shown the ability of microbial communities of different ecosystems to function not only in the presence of bactericidal concentrations of extreme factors, but even in the higher concentrations. The aim of our work was to investigate the quantitative parameters of microbial communities homeostasis in the presence of toxic xenobiotic – NCB. When we talk about homeostasis in this context, we mean the ability of microbial communities to maintain vital functions influenced by extreme factors.

Materials and methods. Our aim was to study the microorganisms homeostasis in the presence of NCB (alien xenobiotic for the biosphere). That is why as objects of study we choose microbial communities of ecosystems differed significantly in physical and chemical factors:

- Mushkarova Yama cave clays ecosystems (Western Ukraine) and Kuybushevskaya (Western Caucasus, Abkhazia) with constant humidity, temperature, organic compounds concentration and absence of lighting;
- mountain jungle (the Andes mountain range, of Papallacta, 4020 m altitude) and the rainforest of Ecuador soils (of Puyo, 950 m altitude), opened ecosystems with dramatically changing climate, temperature and UV radiation.

Museum culture *Escherichia coli* and *Clostridium lituseburense* were selected as reference strains.

Microbial communities quantitative parameters of resistance (CFU, colony morphotypes number and diameter) were determined in liquid and agar media with NCB. As agar media Nutrient Agar (NA HiMedia Ltd.) and oligotrophic agar (OA) were used. As liquid – Nutrient Broth (NB) and oligotrophic broth. In order to get concentration range 50–300 mg/l NCB aliquots of NCB alcohol solution were added in molten and cooled to 45–50 °C agar medium. Petri dishes with NCB agar medium were kept in a desiccator with silicagel for removal of condensation moisture and sterility verification. Then the plates were inoculated with microbial suspensions of samples tenfold dilutions. The agar medium without NCB was used as a sterility

control. Inoculated plates were exposed during 7 days in closed desiccators with a sterile sili-cagel. The number of CFU, colony morphotypes and their diameter were the controlled criteria of microbial communities resistance to NCB.

Strains resistant to NCB were determined in a liquid medium (NB, oligotrophic broth) in a concentration gradient of the xenobiotic (50–300 mg/l). Controlled parameter was the biomass growth, which was determined photocolorimetrically by optical density of the liquid culture on KFK-2MP ($\lambda = 540$ nm). Inoculated medium without xenobiotic was used as a growth control.

Determination of the ability to transform xenobiotic by microbial community and reference strains. Ecuador mountain jungle soil sample (2 g) was cultured in Nutrient Broth (50 ml) in sealed bottles (120 ml) in the presence of xenobiotic (concentration 100 mg/l). Daily museum culture of *E.coli* and *C.lituseburens* were cultured in NB during 10 hours. During the experiment we monitored pH, Eh, optical density, the concentration of xenobiotic and gas phase composition. The optical density was determined on photocolorimeter KFK-2MP ($\lambda = 540$ nm, optical path 0.5 cm). Indexes of pH and Eh were determined using a “pH meter pH-Milivoltmeter MA-150” potentiometer. For Eh measuring we used two electrodes: measuring EVP-1 and chlorosilver comparison electrode EVL-1M3. For pH measuring – combine electrode ESK-10603/4.

Concentration of NCB and chloroaniline (CIA) in the liquid culture was determined by UV absorbance spectrum by using Specord-UV-vis ($\lambda = 200$ –350 nm)¹. Absorption maximum of NCB is 215 nm, CIA – 265 nm. In order to detect xenobiotic concentration NCB was extracted by hexane (1:10) from liquid culture supernatant.

Results and discussion. Samples of Mushkarova Yama and Kuybushevskaya (sampling depth 1 km) caves clays and Ecuador natural tropical ecosystems soils (4020 m height selection and 950 m) were selected in pollution-free areas without anthropogenic and technogenic influence on microbial communities. We determined maximum permissible concentrations of NCB for these ecosystems microbial communities (Fig. 1, 2).

Fig. 1. NCB effect of the CFU number A. Mushkarova Yama; B. Kuybushevskaya

Note: To quantify parameters of resistance of caves microbial communities we used copiocarbotrophic medium (Nutrient Agar – NA with carbon concentration is 850 mg/l) and oligotrophic medium (oligotrophic agar – OA with carbon concentration 85 mg/l) due to the ability of cave microorganisms to grow in a wide range of organic compounds concentrations.

Maximum permissible concentrations of NCB for microbial communities of Mushkarova Yama and Kuybushevskaya clays were 200 and 300 mg/l, respectively. In both cases, the number of surviving microorganisms was greater on the medium with high organic compounds concentrations (NA) comparing with the medium with a low organic compounds concentration – oligotrophic agar (OA).

The maximum permissible concentration of NCB for soil microbial communities of Ecuador mountain jungle and rainforest was 300 mg/l (Fig. 2). It should be noted that in all cases the number of CFU and morphotypes, and the diameter of the colonies decreased while

¹ Absorption in the UV region is determined by the presence of unsaturated bonds in the aromatic ring.

NCB concentration increased. Colony diameter in the concentrations of 100–300 mg NCB/l was reduced in 6–7 times (from 4.5–7 mm to 0.5–2 mm). Morphotypes number decreased from 4–5 to 1–2 in all studied microbial communities.

Fig. 2. NCB effect of the CFU number, Ecuador rainforest and mountain jungle soil samples

Note: To quantify parameters of resistance of tropical microbial communities we used copiocarbotrophic medium (Nutrient Agar - NA, the carbon concentration of 850 mg/l)

Previously we obtained data about the microorganisms resistance in terrestrial ecosystems (Antarctic cliffs, Negev desert soil, black soil Askania Nova, the soil of the Dead Sea, etc.) to extreme factors. It has been shown that there are two types of responses on the effect of extreme factors [2, 3]. The first type of response, also called «correlative», represents inhibition of growth in increasing studied concentration range of xenobiotic. The second type of response represents absence of growth inhibition in the studied xenobiotic concentration range.

Similar types of responses have been shown in studied microbial communities in the NCB concentration gradient. Thus, the correlative response (the first type of response) was characterized for Mushkarova Yama and Ecuador rainforest soil microbial communities. The second type of response was observed for Kuybushevskaya karst cave and Ecuador mountain jungle microbial communities.

Studied microbial communities were not only highly resistant to NCB, but also able to transform it. There are three known ways of nitrochlorinearomatic compounds transformation: nitro group reduction, dehalogenation and aromatic ring reduction [1].

Previously it was shown the non-specific *p*-nitrochlorobenzene reduction on the example of facultative and obligate anaerobic microorganisms museum cultures [5]. Microorganisms can degrade xenobiotic by two ways: reduction to *p*-chloroaniline (*p*-CIA); reduction of the aromatic ring with subsequent cleavage of the cyclic compound.

We investigated the possibility of NCB transforming by Ecuador mountain jungle soil microbial community (Fig. 3).

Fig. 3. NCB transformation by Ecuador mountain jungle soil microbial community

The initial concentration of xenobiotic was 100 mg/l. It decreased in 4 times during cultivation (12 days). The appearance of CIA was not observed. Thus, for Ecuador mountain jungle soil microbial community was shown the second way of NCB destruction – reduction of the aromatic ring with subsequent cleavage of the cyclic compound.

On an example of museum cultures we showed dynamics of NCB reduction (Fig. 3 and Fig. 4). Thus, the growing culture of *E.coli* reduced NCB (initial concentration was 10 mg/l) to CIA in 1.5 hours (Fig.4). A museum culture *C.lituseburens* transformed NCB (160 mg/l) to CIA (60 mg/l) in 10 hours (Fig. 5).

Fig. 4. Transformation of NCB by growing *E.coli* culture

Fig. 5. Transformation of NCB by *Clostridium lituseburens*

In both cases, NCB reduced in CIA independently of the xenobiotic concentration in the medium. The reaction rate increased with a decreasing of redox potential of the medium.

Thus, Ecuador mountain jungle soil microbial community and museum cultures interacted with NCB. However, there were some significant differences. First of all, the museum cultures *E.coli* and *C.lituseburens* transformed NCB in CIA, while Ecuador microbial community provided complete degradation of NCB. This was evidenced by the absence of CIA accumulation in the culture fluid. In addition, in the case of museum cultures decreasing of NCB concentration correlates with decreasing of redox potential. In the experiment with Ecuador microbial community this was not observed. Decreasing of NCB concentration started on the 5th day of cultivation, whereas biomass growth and the change in pH and Eh started on

the 1–2nd day already. So, obviously, during this time Ecuador microbial community adapted to NCB, and only then started interactions that led to the xenobiotic destruction.

Thus, microbial communities of ecosystems with constant physicochemical conditions (cave clays) and Ecuador tropical ecosystems (mountain and rainforest jungle soils) were highly resistant to NCB. They show similar homeostasis parameters in the presence of xenobiotic: the first and the second types of responses on NCB influence, the maximum permissible concentration ranged between 200–300 mg/l. NCB had a pronounced toxic effect on the studied microbial communities. While the NCB concentration increases, CFU, colony number morphotypes and their diameter decreased. The studied microbial communities proceeded complete xenobiotic degradation after adaptation to it.

Таширев О.Б., Суслова О.С., Рокитко П.В.

Институт мікробіології та вірусології ім. Д.К. Заболотного НАН України, Київ

КІЛЬКІСНІ ПОКАЗНИКИ ГОМЕОСТАЗУ МІКРОБНИХ УГРУПОВАНЬ ГРУНТІВ ЕКВАДОРУ ТА ГЛИН КАРСТОВИХ ПОРОЖНИН ЗА ВПЛИВУ П-НІТРОХЛОРБЕНЗОЛУ

Резюме

Показано кількісні показники мікробних угруповань глин карстових порожнин Західної України та Абхазії (Мушкаррова Яма та Куйбишевська) та ґрунтів тропічних екосистем Екватору до дії *n*-нітрохлорбензолу (НХБ). Для зазначених мікробних угруповань встановлено максимально допустимі концентрації та типи відповідей на дію ксенобіотика. Відповідь першого типу характерна для мікробних угруповань глин Мушкаррової Ями та тропічного лісу Екватору. Відповідь другого типу – для мікробних угруповань глин карстової порожнини Куйбишевська та гірських джунглів Екватору. Максимально допустимі концентрації НХБ для Мушкаррової Ями складають 200 мг/л, для інших досліджуваних мікробних угруповань – 300 мг/л. Встановлено, що мікробні угруповання не тільки високостійкі до НХБ, але і взаємодіють із ним, руйнуючи ксенобіотик і знижуючи його концентрацію в 4 рази.

Ключові слова: *n*-нітрохлорбензол, карстові порожнини, тропічні екосистеми Екватору, мікробні угруповання.

Таширев А.Б., Суслова О.С., Рокитко П.В.

Институт микробиологии и вирусологии им. Д.К. Заболотного НАН Украины, Киев

КОЛИЧЕСТВЕННЫЕ ПОКАЗАТЕЛИ ГОМЕОСТАЗА МИКРОБНЫХ СООБЩЕСТВ ПОЧВ ЭКВАДОРА И ГЛИН КАРСТОВЫХ ПОЛОСТЕЙ ПРИ ВЛИЯНИИ П-НИТРОХЛОРБЕНЗОЛА

Резюме

Показаны количественные показатели устойчивости микробных сообществ Западной Украины и Абхазии (Мушкаррова Яма и Куйбышевская) и почв тропических экосистем Экватора к действию *n*-нитрохлорбензола (НХБ). Для указанных микробных сообществ установлены максимально допустимые концентрации НХБ и типы ответов на действие ксенобиотика. Ответ первого типа характерен для микробных сообществ глиин Мушкарровой Ямы и тропического леса Экватора. Ответ второго типа – для микробных сообществ глиин карстовой полости Куйбышевская и горных джунглей Экватора. Максимально допустимые концентрации НХБ для Мушкарровой Ямы составляют 200 мг/л, для других исследуемых микробных сообществ – 300 мг/л. Установлено, что микробные сообщества не только высокоустойчивы к НХБ, но и взаимодействуют с ним разрушая ксенобиотик и снижая его концентрацию в 4 раза.

Ключевые слова: *n*-нитрохлорбензол, карстовые полости, тропические экосистемы Экватора, микробные сообщества.

1. *Gvozdyak P.I., Mogilevich N.F., Tashyrev A.B.* Transformation of *p*-nitrochlorobenzene by *E.coli* // *Microbiologiya*. – 1983. – Vol. 52. – № 1. – P. 22-25.
2. *Tashyrev A.B., Matveeva N.A., Romanovskaya V.A., Tashyreva A.A., Rokitko P.V.* Polyresistance superstability and heavy metals Antarctic microorganisms // *Dopovidi Nacional'noi Akademii Nauk Ukraïni*. – 2007. – № 11. – P. 170-175.
3. *Tashyrev A.B., Rokitko P.V., Levishko A.S., Romanovskaya V.A., Tashyreva A.A.* Resistance to toxic metals of chemoorganotrophic bacteria isolated from the Antarctic cliffs // *Mikrobiol. zhurnal*. – 2012. – 74, № 2. – P. 3-7.
4. *Tashyrev A.B., Romanovskaya V.A., Rokitko P.V., Tashyreva A.A.* Multiple resistance to toxic metals of Antarctic cliffs microorganisms (o.Galindez) // *Ukr. antarktichny zhurnal*. – 2011-2012. – № 11-12. – P. 212-221.
5. *Tashyrev A.B., Tihnenko S.A.* Transformation of *p*-nitrochlorobenzene by obligate anaerobic microorganisms. – In the book. “Microbiology of water treatment”: Abstracts 2 Union Conference. Kiev, 1982, p.194.
6. *Li Haixiang, Yang Xin, Zhong Fohua, Li Junying, Xia Siqing.* Experimental study on para-nitrochlorobenzene (p-NCB) removal from groundwater by autohydrogenotrophic microorganism// *Chinese Journal of Environmental Engineering*. – 2011. – № 2. – P. 43-48.

Отримано 10.06.2014